
Physics 12 Booklet #4

ı	n	d	п	c	ti	o	n
		ч	u	•			

It was discovered that a wire with current flowing the was: Could a	rough it created a r cause <b>CUN</b>	magnetic f	field, the next question to flow in a wire?
Another way to say this is: Could a magnetic field ge	nerate anE^	1F	?
Michael Faraday discovered it was possible with a			
To determine the direction of the induced current w		in	the

direction which opposes the change causing it.





In order to calculate the EMF generated we need to use the idea of magnetic flux.


Magnetic Flux

Possing through a coil

More flux

Less flux

Symbol 1



0.0Um

Magnetic flux perpendicular to a magnetic field can be calculated as

Example: A square loop of wire is perpendicular to a 1.50 T magnetic field. If each side of the wire is

2.10 cm what is the magnetic flux through the loop?

 $A = (0.021)^2 = 0.000441m^2$ 

$$\Phi = (1.5)(0.000441) = 0.000662 Wb$$

The EMF is induced on a loop of wire by a changing flux

$$\mathcal{E} = \left| \frac{\Delta \Phi}{\tau} \right|$$

If there are several loops of wire the EMF generated by each loop adds together so total EMF is

Example: A circular loop of wire with radius 2.5 cm is placed in a magnetic field B=0.020 T into the page. The field is reduced to 0.010 T into the page in 0.10 seconds. Aced = Tr = Tr (0.025)

a) What is the average induced EMF?

= 0.001963m

$$\tilde{\zeta}(\tilde{\zeta})^* \to \tilde{\zeta}_{\chi}$$

b) Which direction does the current flow?

Change Less into Oppose: More into Clackwise

$$\Delta \Phi = \Phi_{\phi} - \Phi_{\phi};$$

$$= (0.001963)(0.010) - (0.001963)(0.020)$$

$$= -1.96 \times 10^{-5} \text{ Wb}$$

$$E = \left| \left( \frac{-1.96 \times 10^{-5} \text{ Wb}}{0.10} \right) \right|$$

$$= 1.96 \times 10^{-4} \text{ V} \quad ^{30}$$